

ON c -NORMAL AND S -QUASINORMALLY EMBEDDED SUBGROUPS OF A FINITE GROUP

Hong Pan

*College of Mathematics and Statistics, Xinyang Normal University,
Xinyang 464000, People's Republic of China
fz1105678@163.com*

Abstract. If P is a p -group for some prime p we shall write $\mathcal{M}(P)$ to denote the set of all maximal subgroups of P and $\mathcal{M}_d(P) = \{P_1, \dots, P_d\}$ to denote any set of maximal subgroups of P such that $\bigcap_{i=1}^d P_i = \Phi(P)$ and d is as small as possible. In this paper, the structure of a finite group G under some assumptions on the c -normal or S -quasinormally embedded subgroups in $\mathcal{M}_d(P)$, for each prime p , and Sylow p -subgroups P of G is researched. Some known results are generalized.

Keywords: c -normal subgroup, S -quasinormally embedded subgroup, supersolvable groups.

1. Introduction

All groups considered in this paper are finite. Let G be a group and let $\mathcal{M}(G)$ be the set of all maximal subgroups of all Sylow subgroups of G . A interesting topic in group theory is to study the influence of the elements of $\mathcal{M}(G)$ on the structure of G . A typical result in this direction is due to Srinivasan [1]. He proved that G is supersolvable provided that every member of $\mathcal{M}(G)$ is normal in G . This result has been widely generalized.

A subgroup H of G is called s -quasinormal in G provided H permutes with all Sylow subgroups of G , i.e, $HP = PH$ for any Sylow subgroup P of G . This concept was introduced by Kegel in [2] and has been studied extensively by Deskins [3] and Schmidt [4]. More recently, Ballester-Bolinches and Pedraza-Aguilera [5] generalized s -quasinormal subgroups to s -quasinormally embedded subgroups. A subgroup H of G is said to be s -quasinormally embedded in G provided every Sylow subgroup of H is a Sylow subgroup of some s -quasinormal subgroup of G . In [5], Ballester-Bolinches and Pedraza-Aguilera showed that, if every subgroup in $\mathcal{M}(G)$ is s -quasinormally embedded in G , then G is supersolvable. Assad and Heliel [6] showed that G is p -nilpotent for the smallest prime p dividing $|G|$ if and only if all members of $\mathcal{M}(P)$ are s -quasinormally embedded in G , where P is a Sylow p -subgroup of G .

As another generalization of the normality, Wang [7] introduced the following concept: A subgroup H of G is called c -normal in G if there is a normal subgroup K such that $G = HK$ and $H \cap K \leq H_G$, where H_G is the normal core of H in G . In [7], Wang showed that G is supersolvable if every member of $\mathcal{M}(G)$

is c -normal. Wang's result has been generalized by some authors (see [8-12], etc). For example, Guo and Shum showed in [8] the following result. Let p be the smallest prime dividing the order of G and let P be a Sylow p -subgroup of G . If every member of $\mathcal{M}(P)$ is c -normal, then G is p -nilpotent. The research on c -normal subgroups has formed a series, which is similar to the series of s -quasinormal subgroups. However, the two series are independent of each other. The aim of this article is to unify and improve the results of [1], [5], [7] and some of [8].

If P is a p -group for some prime p we shall write $\mathcal{M}(P)$ to denote the set of all maximal subgroups of P and $\mathcal{M}_d(P) = \{P_1, \dots, P_d\}$ to denote any set of maximal subgroups of P such that $\bigcap_{i=1}^d P_i = \Phi(P)$ and d is as small as possible.

Such subset $\mathcal{M}_d(P)$ is not unique for a fixed P in general. We know that

$$|\mathcal{M}(P)| = \frac{p^d - 1}{p - 1}, |\mathcal{M}_d(P)| = d, \lim_{d \rightarrow \infty} \frac{p^d - 1}{(p - 1)d} = \infty,$$

so $|\mathcal{M}(P)| \gg |\mathcal{M}_d(P)|$.

In this paper, we study the influence of the members of some fixed $\mathcal{M}_d(P)$ on the structure of group G . Our results are more general.

2. Basic definitions and preliminary results

In this section, we give some results that are needed in this paper.

Lemma 2.1 ([5]). *Suppose that U is an s -quasinormally embedded subgroup of G and that K is a normal subgroup of G . Then:*

- (a) U is s -quasinormally embedded in H whenever $U \leq H \leq G$;
- (b) UK is s -quasinormally embedded in G and UK/K is s -quasinormally embedded in G/K .

Lemma 2.2 ([7]). *Let $X \leq H \leq G$ and $N \trianglelefteq G$. Then:*

- (a) If X is c -normal in G , then X is also c -normal in H ;
- (b) If X is c -normal in G , then XN/N is c -normal in G/N .

Lemma 2.3 ([3]). *If H is an s -quasinormal subgroup of the group G , then H/H_G is nilpotent.*

Lemma 2.4 ([4]). *For a nilpotent subgroup H of G , the following two statements are equivalent:*

- (a) H is s -quasinormal in G ;
- (b) The Sylow subgroups of H are s -quasinormal in G .

Lemma 2.5 ([6]). *Let G be a group and let P_0 be a maximal subgroup of P . Then the following two statements are equivalent:*

- (a) P_0 is normal in G ;
- (b) P_0 is s -quasinormal in G .

Lemma 2.6 ([13]). *If P is a Sylow p -subgroup of G and $N \trianglelefteq G$ such that $P \cap N \leq \Phi(P)$, then N is p -nilpotent.*

Lemma 2.7 ([14]). *A group G is supersolvable if and only if there exists a subgroup of order dividing $|H|$ for every subgroup H of G .*

Lemma 2.8 ([15]). *Let p_1 be the minimal prime dividing $|G|$ and p_s the maximal prime dividing $|G|$. If G possesses two supersolvable subgroups H and K with $|G : H| = p_1$ and $|G : K| = p_s$, then G is supersolvable.*

Lemma 2.9 ([16]). *Let H be a p -subgroup of G . Then, the following statements are equivalent:*

- (a) H is s -quasinormal in G ;
- (b) $H \leq O_p(G)$, and H is s -quasinormally embedded in G .

3. Main results

Theorem 3.1. *Let p be the smallest prime dividing the order of a group G and P be a Sylow p -subgroup of G . Assume that every member of some fixed $\mathcal{M}_d(P)$ is either c -normal or s -quasinormally embedded in G . Then G is p -nilpotent.*

Proof. Assume that the result is not true and let G be a counterexample of minimal order. Let $\mathcal{M}_d(P) = \{P_1, \dots, P_d\}$. By hypothesis, each P_i is either c -normal or s -quasinormally embedded in G . Without loss of generality, let I_1 be the subset of $\{1, \dots, d\}$ such that every $P_i (i \in I_1)$ is c -normal in G and I_2 is the subset such that every $P_i (i \in I_2)$ is s -quasinormally embedded in G . We prove the theorem by the following claims:

- (1) $O_{p'}(G) = 1$.

Set $N = O_{p'}(G)$. Consider the quotient group G/N . We know that PN/N is a Sylow p -subgroup of G/N , $N_{G/N}(PN/N) = N_G(P)N/N$ and $\mathcal{M}(PN/N) = \{P_1N/N, \dots, P_mN/N\}$. Now, by Lemma 2.1 and Lemma 2.2, we see easily that G/N satisfies the condition. If $O_{p'}(G) > 1$, then $G/O_{p'}(G)$ is p -nilpotent and hence G itself is p -nilpotent, a contradiction. Thus claim (1) holds.

- (2) G/P_{iG} is p -nilpotent for all $i \in I_1$, where P_{iG} is the core of P_i in G .

In this case, P_i is a c -normal subgroup of G . We know that there exists a normal subgroup K_i of G such that $G = P_iK_i$ and $P_i \cap K_i = P_{iG}$. Hence,

$$G/P_{iG} = P_i/P_{iG} \cdot K_i/P_{iG}, P_i \cap K_i = P_{iG}.$$

Therefore,

$$|K_i/P_{iG}|_p = |G : P_i|_p = |P : P_i| = p.$$

As p is the smallest prime dividing $|G|$, we know that K_i/P_{iG} is p -nilpotent by Burnside's theorem. Therefore, K_i/P_{iG} has a normal Hall p' -subgroup H/P_{iG} . We see that H/P_{iG} is also a normal Hall p' -subgroup of G/P_{iG} because K_i/P_{iG} is normal in G/P_{iG} . It follows that G/P_{iG} is p -nilpotent for all $i \in I_1$.

For every $P_i (i \in I_2)$, there exists an S -quasinormal subgroup H_i of G such that P_i is a Sylow p -subgroup of H_i .

(3) G/H_{iG} is p -nilpotent for all $i \in I_2$, where H_{iG} is the core of H_i in G .

In fact, As H_i is an S -quasinormal subgroup of G and P_i is a Sylow p -subgroup of H_i , it follows that H_i/H_{iG} is S -quasinormal in G/H_{iG} , and the Lemma of 2.3 asserts that H_i/H_{iG} is nilpotent. Hence, H_i/H_{iG} is an S -quasinormal nilpotent subgroup of G/H_{iG} . By Lemma 2.4, every Sylow subgroup of H_i/H_{iG} is S -quasinormal in G/H_{iG} . Since $P_i H_i/H_{iG}$ is a Sylow p -subgroup of H_i/H_{iG} , it follows that $P_i H_i/H_{iG}$ is S -quasinormal in G/H_{iG} . Thus, Lemma 2.5 indicates that $P_i H_i/H_{iG}$ is normal in G/H_{iG} . Therefore, $P_i H_{iG} \trianglelefteq G$. Noting that P_i is a Sylow p -subgroup of H_i , we have $P_i \leq H_{iG}$. Therefore, $|G/H_{iG}|_p = p$. Now, as p is the smallest prime dividing $|G|$, by Burnside's theorem, we see that G/H_{iG} is p -nilpotent for each $i \in I_2$, which proves (3).

Let

$$N = \left(\bigcap_{i \in I_1} P_{iG} \right) \cap \left(\bigcap_{i \in I_2} H_{iG} \right).$$

(4) N is p -nilpotent.

First, as all P_{iG} and H_{iG} are normal in G , we get $N \trianglelefteq G$. Second, we consider the subgroup $P \cap N$. Recall that P_i is a Sylow p -subgroup of H_{iG} and $P_i \leq P$, so $P \cap H_{iG} \leq P_i$. Moreover, $P_i \leq P \cap H_{iG}$. We have $P \cap H_{iG} = P_i$. Therefore,

$$P \cap N = \left(\bigcap_{i \in I_1} P_{iG} \right) \cap \left(\bigcap_{i \in I_2} H_{iG} \cap P \right) = \left(\bigcap_{i \in I_1} P_{iG} \right) \cap \left(\bigcap_{i \in I_2} P_i \right) = \Phi(P).$$

Applying Lemma 2.6, we know that N is p -nilpotent.

(5) Final contradiction.

Now, N possesses a Hall p' -normal subgroup $N_{p'}$ such that $N = N_p N_{p'}$, where N_p is a Sylow p -subgroup of N . Then, $N_{p'} \text{ char } N \trianglelefteq G$, so $N_{p'}$ is normal in G , and hence, $N_{p'} \leq O_{p'}(G)$. It follows by $O_{p'}(G) = 1$ that $N_{p'} = 1$. Consequently, N is a normal p -subgroup of G , and so, $N = P \cap N = \Phi(P)$. Also, note that the class of p -nilpotent groups is a formation, by steps (2) and (3), we have G/N must be p -nilpotent. It follows that $G/\Phi(P)$ is p -nilpotent. Moreover, by III, 3.3 Hils-Satz in [17], $\Phi(P) \leq \Phi(G)$, so $G/\Phi(G)$ is p -nilpotent. It follows that G would be p -nilpotent, contrary to the choice of G .

The following corollaries are immediate from Theorem 3.1.

Corollary 3.2 ([8]). *Let p be the smallest prime dividing the order of G and let P be a Sylow p -subgroup of G . If every member of $\mathcal{M}(P)$ is c -normal, then G is p -nilpotent.*

Corollary 3.3 ([6]). *Let p be the smallest prime dividing the order of G and let P be a Sylow p -subgroup of G . If every member of $\mathcal{M}(P)$ is s -quasinormally embedded in G , then G is p -nilpotent.*

Theorem 3.4. *Let G be a group. If there exists a normal subgroup H of G such that G/H is supersolvable, and for each Sylow subgroup P of H , every member in some fixed $\mathcal{M}_d(P)$ is either c -normal or s -quasinormally embedded in G , then G is supersolvable.*

Proof. Suppose that the theorem is false so that there exists a counterexample G of minimal order. We shall finish the proof by the following claims.

(1) H is a q -group for some prime q .

By hypothesis and Theorem 3.1, we have that H has a Sylow tower of supersolvable type. Let q be the largest prime dividing $|H|$, and let Q be a Sylow q -subgroup of H . The property that H possesses an order Sylow tower property implies that Q is normal in H . Now, $Q \text{ char } H$ and $H \trianglelefteq G$, so $Q \trianglelefteq G$. Furthermore, $(G/Q)/(H/Q) \cong G/H$, and Lemmas 2.1 and 2.2 show that G/Q satisfies the condition of the theorem, by the choice of G , G/Q is supersolvable. Hence, $H = Q$ by the choice of H .

(2) Q is a Sylow q -subgroup of G .

Suppose that Q is not a Sylow q -subgroup of G . Let p be the smallest prime dividing $|G/Q|$ and r the largest prime dividing $|G/Q|$. By (1), G/Q is supersolvable. By Lemma 2.7, G/Q contains two subgroups M_1/Q and M_2/Q with $|G : M_1| = p$ and $|G : M_2| = r$. By Lemmas 2.1 and 2.2, (M_i, Q) ($i = 1, 2$) satisfy the condition of the theorem. By the choice of G , M_1 and M_2 are supersolvable. Now, by Lemma 2.8, G would be supersolvable, which is a contradiction. Thus, (2) holds.

(3) $\Phi(Q) = 1$.

Otherwise, by Lemmas 2.1 and 2.2, $G/\Phi(Q)$ satisfies the hypothesis, applying induction, we have $G/\Phi(Q)$ is supersolvable. Furthermore, $\Phi(Q) \leq \Phi(G)$ by III, 3.3 Hilfs-satz in [17], so $G/\Phi(G)$ is supersolvable. It follows that G is supersolvable, which is a contradiction.

(4) Q is a minimal normal subgroup of G .

Let N be a minimal normal subgroup of G contained in Q . Clearly the quotient group $(G/N, Q/N)$ satisfies the condition, so G/N is supersolvable. As the class of supersolvable groups is a formation, N must be the unique minimal normal subgroup of G which is contained in Q and $N \not\leq (G)$. So there is a maximal subgroup M of G such that $G = NM$ and $N \cap M = 1$. Thus $Q = N(Q \cap M)$. As $G = QM$ and Q is normal abelian in G , we know that $Q \cap M$ is normal in G . If $Q \cap M > 1$, let N_1 be a minimal normal subgroup of G such that $N_1 \leq Q \cap M$, then $N_1 \leq Q$ and $N \neq N_1$, this is a contradiction. Hence $Q \cap M = 1$, which implies $Q = N$.

(5) Every $Q_i \in \mathcal{M}_d(Q) = \{Q_1, \dots, Q_d\}$ is S -quasinormally embedded in G .

Assume that there is a Q_i in $\mathcal{M}_d(Q)$ such that Q_i is c -normal in G . By definition, there is a normal subgroup K_i of G such that $G = Q_i K_i$ and $Q_i \cap K_i = Q_{iG}$ is a normal subgroup of G . By (4), $Q_i \cap K_i = 1$ or Q . If $Q_i \cap K_i = Q$, then $Q_i = Q$, a contradiction. If $Q_i \cap K_i = 1$, then $Q = Q_i(Q \cap K_i)$. But then $Q \cap K_i$ is a normal subgroup of order q of G . So $Q = Q \cap K_i$ by (4). As the class of supersolvable groups is a formation, thus G is supersolvable, contrary to the choice of G .

(6) $Q_i (i = 1, 2, \dots, d)$ are normal subgroups of G .

Lemmas 2.5 and 2.9 imply that $Q_i (i = 1, 2, \dots, d)$ are normal subgroups of G .

(7) The final contradiction.

Now,

$$(G/Q_i)/(Q/Q_i) \cong G/Q,$$

by (1), G/Q is supersolvable. As Q/Q_i is cyclic of order q , it follows that G/Q_i is supersolvable. Set

$$N = \bigcap_{i=1}^d Q_i.$$

By the definition of $\mathcal{M}_d(Q)$,

$$\bigcap_{i=1}^d Q_i = \Phi(Q),$$

so $N = \Phi(Q)$. Now, by the class of supersolvable groups is a formation, $G/\Phi(Q)$ is supersolvable. It follows that $G/\Phi(G)$ is supersolvable, and hence, G is supersolvable. which is a final contradiction. The proof is now completed.

The following corollaries are immediate from Theorem 3.4.

Corollary 3.5 ([7]). *Let G be a group. If every member of $\mathcal{M}(G)$ is c -normal, then G is supersolvable.*

Corollary 3.6 ([5]). *Let G be a group. If every subgroup in $\mathcal{M}(G)$ is s -quasinormally embedded in G , then G is supersolvable.*

References

- [1] S. Srinivasan, *Two sufficient conditions for supersolvability of finite groups*, Israel J. Math., 35(1980), 210-214.
- [2] O.H. Kegel, *Sylow Gruppen und subnormalteiler endlicher Gruppen*, Math. Z., 78 (1962), 205-221.
- [3] W.E. Deskins, *On quasinormal subgroups of finite groups*, Math. Z., 82 (1963), 125-132.

- [4] P. Schmid, Subgroups permutable with all Sylow subgroups, *J. Algebra* 207(1998), 285-293.
- [5] A. Ballester-Bolínches and M.C. Pedraza-Aguilera, *Sufficient conditions for supersolvability of finite groups*, *J. Pure Appl. Algebra*, 127 (1998), 113-118.
- [6] M. Asaad and A.A. Heliel, *On S -quasinormal embedded subgroups of finite groups*, *J. Pure Appl. Algebra*, 165 (2001), 129-135.
- [7] Yanming Wang, *c -normality of groups and its properties*, *J. Algebra*, 180 (1996), 954-965.
- [8] X.Y. Guo and K. P. Shum, *On c -normal maximal and minimal subgroups of Sylow p -subgroups of finite groups*, *Arch. Math.*, 80 (2003), 561-569.
- [9] D. Li and X. Guo, *The influence of c -normality of subgroups on structure of finite groups*, *Comm. Algebra*, 26 (1998), 1913-1922.
- [10] D. Li and X. Guo, *The influence of c -normality of subgroups on structure of finite groups II*, *J. Pure Appl. Algebra*, 150 (2000), 53-60.
- [11] H. Wei, *On c -normal maximal and minimal subgroups of Sylow subgroups of finite groups*, *Comm. Algebra*, 29 (2001), 2193-2200.
- [12] H. Wei, Y. Wang and Y. Li, *On c -normal maximal and minimal subgroups of Sylow subgroups of finite groups II*, *Comm. Algebra*, 31 (2003), 4807-4816.
- [13] J. Tate, *Nilpotent quotient groups*, *Topology*, 3 (1964), 109-111.
- [14] M. Weinstein, *Between Nilpotent and Solvable*, Passaic: Polygonal Publishing House, 1982.
- [15] M. Asaad, *On the supersolvability of finite groups*, *Annales Univ Sci, Budapest*, XIII (18) (1975), 3-7.
- [16] Li Shirong, Shen Zhencai, Liu Jianjun, Liu Xiaochun, *The influence of SS -quasinormality of some subgroups on the structure of finite groups*, *J. Algebra*, 319 (2008), 4275-4287.
- [17] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin-Heidelberg-New York, 1967.

Accepted: 16.08.2016